



Predicting safe sandwich production

Birk, Tina; Duan, Zhi; Møller, Cleide Oliveira de Almeida; Hansen, Tina Beck; Friis Hansen, Heidi; Knøchel, Susanne

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Birk, T., Duan, Z., Møller, C. O. D. A., Hansen, T. B., Friis Hansen, H., & Knøchel, S. (2014). *Predicting safe sandwich production*. Poster session presented at The Danish Microbiological Society Annual Congress 2014, Copenhagen, Denmark.


General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

DTU¹ Predicting safe sandwich production



Tina Birk¹, Zhi Duan^{1,2}, Cleide Oliveira de Almeida Møller¹, Heidi Friis Hansen³, Susanne Knøchel² and Tina Beck Hansen¹






Have you ever experienced that 3 hours was not enough time to prepare, distribute and serve ready-to-eat sandwiches? If yes, this decision support tool might be for you. The tool helps you to control sandwich production by predicting growth of foodborne pathogens.

Using time/temperature measurements obtained during preparation, in combination with information on prehistory of the ingredients as well as the expected time/temperature conditions of distribution and serving, the growth of *Listeria monocytogenes*, *Salmonella* and psychrotrophic *Clostridium botulinum* can be predicted. Based on these predictions, the tool determines whether any of the lag times have been exceeded during the total preparation, distribution and serving time. All underlying growth models use a "worst case" ingredient identified as cooked, sliced chicken.


Danish hygiene guidelines (nr. 9025 af 17/01/2013, afsnit 26.3)
Violation of the cold chain

- As per the guidelines, chilled foods should not be outside the cold chain for more than 3 hours, including the time for preparation and serving.
- A prolongation of the acceptable time outside the cold chain can be allowed if it can be documented that the prolongation does not result in any health risks.


Ex. growth curves




Constant conditions



Lag time may not be exceeded
lag time = RLT · generation time
RLT: relative lag time is a constant




Dynamic conditions



$$N(t) = \sum \frac{\ln(2)}{(b \cdot (T - T_{\min}))^2} \cdot \frac{1}{\Delta t}$$

where b and T_{\min} are constants, specific for each pathogen

SAFE, if $N(t) < \text{RLT}$



For further details contact Tina Beck Hansen, email: tibha@food.dtu.dk

